

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 01 JUL 2003		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Multiplexed Pathogenicity Island Detection Of Biowarfare Agents, Environmental Pathogens, And Food-Borne Pathogens				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Institute for Advanced Technology The University of Texas at Austin Austin, Texas 78759				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM001523.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

MULTIPLEXED PATHOGENICITY ISLAND DETECTION OF BIOWARFARE AGENTS, ENVIRONMENTAL PATHOGENS, AND FOOD-BORNE PATHOGENS

Robert Chin*, Steve Kornguth, Shelley Payne
Institute for Advanced Technology
The University of Texas at Austin
Austin, Texas 78759

James J. Valdes
The US Army SBCCOM
Aberdeen Proving Ground, MD 21010

Kerry Oliver
Radix BioSolutions, Ltd.
Georgetown, Texas 78626

ABSTRACT

The Institute for Advanced Technology (IAT) at The University of Texas at Austin (UT Austin) and UT faculty have joined together with Radix BioSolutions, Ltd. (Georgetown, TX) to transition technologies developed at UT Austin. The long-term objective of this project is to provide a fieldable system composed of a sampler and pre-sample preparation module and a transduction platform. The envisioned system combines pathogenicity island sequences with a multiplexed assay platform to develop a rapid, extensive infectious pathogen sensor. This sensor is based upon the Luminex xMAP™ System, a multiplexed assay platform that combines high sample throughput (up to 600 samples/hour) with high information content per sample (up to 100 parameters tested simultaneously per sample). Pathogenicity islands are DNA sequences that directly relate the infectious properties of an organism to its host. By quickly screening for multiple pathogenicity island sequences, end-users will have the capability to detect the first signs of a bioattack without requiring screening for a particular organism. Once a pathogenicity “fingerprint” is identified, an alert is transmitted to an end user, which includes military, health providers, and other government agencies.

INTRODUCTION

Recent events in the U.S. have validated the fear that terrorists will use biowarfare agents in the homeland (anthrax) or on U.S. troops abroad (Congo-Crimean Hemorrhagic fever). Present biological agent detection systems tend to rely on the detection of single, or a few, DNA sequences, antigens or antibodies. The objective for this work is to use identified pathogenicity islands to completely define the critical domains of a set of virulence genes of key pathogens, and to use this information to design and build a prototype multiplexed detection system that will rapidly screen and detect every virulent mutant of that pathogen or pathogen class, and will give no false alarms. This detection system will have hundreds of probes (multiplexed), so that false positives and false negatives will be determined immediately. The system will be rugged, lightweight, and low-cost. A side benefit of this type of multiplexed detection is that it may inform investigators if a particular pathogen is man-made, if it is a new emergent infectious disease, or the source of the strain. The enteric pathogens include a large number of related, but distinct organisms. Identification of the organisms has been based on antigenic or biochemical analysis. While

this approach is effective, it may fail to identify potential pathogens if they are antigenic variants or differ in metabolic pathways. For example, epidemics of cholera have traditionally been caused by *Vibrio cholerae* serotype O1 and antisera against the O1 antigen was used to detect epidemic strains. However, a previously unrecognized epidemic serotype, O139, that was not detected by O1 antisera emerged recently and caused a major pandemic of cholera. An alternative approach to rapid detection and identification of pathogens is to develop methods that detect gene encoding virulence factors associated with a particular syndrome rather than detecting bacteria that may or may not have these virulence factors. We have applied this approach to rapid identification of pathogens in the *E. coli/Shigella* group. These pathogens contain the same “core” sequences but have additional regions of DNA encoding virulence determinants that appear to have been inserted into the chromosomes (as pathogenicity islands and prophages) or are carried on extrachromosomal elements (plasmids). It is the particular set of virulence factors encoded by a member of this group that determines pathogenic potential. *E. coli* O157:H7, which has been associated with large outbreaks of hemorrhagic colitis and hemolytic uremic syndrome has the basic *E. coli* K-12 genome but also carries a phage encoding the shiga-like toxin, the

LEE pathogenicity island carrying the attachment and effacing loci, the *Shu* heme transport island, and plasmids encoding virulence factors. *Shigella dysenteriae*, which causes dysentery, encodes a related Shiga toxin and has the same *Shu* island, but lacks the *LEE* and has a different plasmid that confers the ability to invade and multiply within human epithelial cells (Sansonetti 1992, Wyckoff et al. 1998). *S. flexneri* lacks the Shiga toxin locus and the *Shu* island (Wyckoff et al. 1998) but carries an island encoding a different iron transport system (aerobactin) that is not found in *S. dysenteriae* and *E. coli* O157:H7 (Vokes et al. 1999). By identifying the specific factors present in a member of this group, we can determine its potential to cause disease and the type of disease. This could lead to more rapid identification of the pathogen and determination of the proper course of treatment.

The Luminex xMAP™ System provides a rapid and inexpensive method to simultaneously detect the presence of antigens or specific DNA sequences. The xMAP™ System utilizes polystyrene microspheres internally dyed with two spectrally distinct fluorochromes. Using precise ratios of these fluorochromes, a suspension array is created consisting of 100 spectrally distinct microsphere sets. Each microsphere set can possess a different reactant on its surface, and thus, up to 100 different analyses can be detected simultaneously in a single reaction vessel. A third fluorochrome coupled to a reporter molecule quantifies the biomolecular interaction that has occurred at the microsphere surface. The multiplexing capabilities of the xMAP™ system provide an assay detection platform that does not require users to screen for specific organisms, but allows a more thorough screen for the presence of biowarfare, food-borne, and environmental pathogens. As new pathogenicity islands are discovered or mutations are determined, new microsphere sets carrying these sequences will be added to the current assay kit, thus seamlessly updating the kit.

PRELIMINARY RESULTS

Recent experiments have shown that the instrument is capable of detecting multiple pathogens at less than 1000 genome copies per amplified sample. An initial pathogenicity island multiplexed assay has been developed that examines three different sequences, two for pathogenicity islands and one for a common junction area insertion site. Preliminary data demonstrates a 3.5-7 fold signal to noise ratio using as little as 1µL of PCR product for detection (Figures below).

CONCLUSION

The use of pathogenicity islands can be used as indiscriminant identifiers of pathogens thus providing a genomic fingerprint to identify potential of pathogens. The use of the Luminex xMAP™ system provides for a rapid high throughput screening system which can be modified by adding additional gene sequences as need to current sequences or immunoassays. Because the detection system is based on

multiple sequences for each pathogenicity factor one can essentially eliminate false positives and false negatives.

The proposed system has commercial applications in office health monitoring, hospital infectious disease detection, environmental and agricultural pathogen monitoring, and food-borne pathogen monitoring.

REFERENCES

- Purdy, G. and Payne, S.M., "The SHI-3 Iron Transport Island of *Shigella boydii* 0-1392 Encodes the Genes for Aerobactin Synthesis and Transport," *J. Bacteriol.*, Vol. 183, pp. 4176-4182, 2001.
- Sansonetti, P.J., "Molecular and Cellular Biology of *Shigella flexneri* Invasiveness: From Cell Assay Systems to Shigellosis," *Curr Top Microbiol. Immunol.*, Vol. 180, pp. 1-19, 1992.
- Vokes, S., Reeves, S. A., Torres, A. G., and Payne, S. M., "The Aerobactin Iron Transport System Genes in *Shigella flexneri* are Present within a Pathogenicity Island," *Mol. Microbiol.*, Vol. 33, pp. 63-73, 1999.
- Wyckoff, E.E., Duncan, D., Torres, A., Maase, K., and Payne, S.M., "Structure of the *Shigella dysenteriae* Haem Transport Locus and its Phylogenetic Distribution in Enteric Bacteria," *Mol. Microbiol.*, Vol. 28, pp. 1139-1152, 1998.

ACKNOWLEDGEMENTS

The research reported in this presentation was performed in connection with Contract number DAAD17-01-D-0001 with the U.S. Army Research Laboratory. The views and conclusions contained within this presentation are those of the authors and should not be interpreted as presenting the official policies or position, either expressed or implied, of the U.S. Army Research Laboratory or the U.S. Government unless so designated by other authorized documents. Citation of manufacturer's or trade names does not constitute an official endorsement or approval of the use thereof. The U.S. Government is authorized to reproduce and distribute reprints for Government purposes notwithstanding any copyright notation hereon.